



BCL6 inhibitor | BI-3812

Table of contents

Summary	2
Chemical Structure	2
Highlights	3
Target information	3
<i>In vitro</i> activity	4
<i>In vitro</i> DMPK and CMC parameters	4
Negative control	5
Selectivity	5
Co-crystal structure of the BI probe compound and the target protein Reference molecule(s)	5
Several small molecule BCL6 inhibitors have been published recently ^{7,8,9,10} .	5
Summary	6
References	6

Summary

BI-3812 is a single digit nanomolar BCL6::Co-repressor inhibitor which also inhibits the BCL6::Co-repressor complex formation in cells.

Chemical Structure




Figure 1: 2-D structure of BI-3812, a BCL6 inhibitor




Figure 2: BI-3812, 3D conformation, based on X-ray structure with BI-3802

Highlights

B-cell lymphoma 6 (BCL6) is a known oncogenic driver and frequently overexpressed in many DLBCL. BI-3812 potently inhibits the interaction of the BTB/POZ domain of BCL6 with several co-repressors *in vitro* ($IC_{50} \leq 3$ nM). In a cellular context, BI-3812 inhibits the BCL6::Co-repressor complex formation with an IC_{50} of 40 nM¹. The high potency and good permeability properties of BI-3812 make this molecule a very good cellular probe compound for testing hypotheses around BCL6 biology. With BI-5273 we also offer a structurally close analog which can be used as a negative control for *in vitro* experiments ($IC_{50} \sim 10$ μ M)¹.

Target information

B-cell lymphoma 6 (BCL6) functions as a transcriptional repressor that binds specific DNA sequences *via* its Zn-fingers and recruits transcriptional co-repressors (e.g. BCOR, SMRT, NCOR) by its BTB/POZ domain². BCL6 is essential for the germinal center (GC) reaction³. It represses a broad set of genes that are required to sustain mutagenic activity without activating the DNA damage response or apoptosis⁴. BCL6 also prevents maturation to plasma or memory cells and helps to maintain a de-differentiated state. Its expression must be switched off to allow the B-cell to exit the GC cycle and differentiate. BCL6 is a known oncogenic driver of DLBCL^{5,6} and frequently overexpressed in DLBCL.




Figure 3: BCL6-BTB dimer with BI-3802, as observed by X-ray.¹ BI-3802, a close analog of BI-3812, binds at the interface of two monomers (monomers are shown in green and grey).

***In vitro* activity**

BI-3812 displays an $IC_{50} \leq 3$ nM in a BCL6::BCOR *ULight* TR-FRET assay.

PROBE NAME / NEGATIVE CONTROL	BI-3812	BI-5273
MW [Da]	558	500
BCL6::BCOR <i>ULight</i> TR-FRET (IC_{50}) [nM] ^a	≤ 3	10,162
BCL6::NCOR LUMIER (IC_{50}) [nM]	40	n.d.

^aWith affinities of approximately 3 nM, the assay wall of this assay is reached, limiting the accuracy of the biochemical assay.

We recommend to store and use 1 mM DMSO stock solutions of BI-3812 for all *in vitro* experiments.

***In vitro* DMPK and CMC parameters**

PROBE NAME / NEGATIVE CONTROL	BI-3812	BI-5273
Aqueous solubility @ pH 6.8 [μ g/ml]	< 1	84
CACO permeability @ pH 7.4 [$*10^{-6}$ cm/s]	2.8	22
CACO efflux ratio	14	0.6
Human hepatocyte clearance [% Q_H]	n.d.	n.d.
Plasma protein binding human [%]	96.89	n.d.

Negative control

BI-5273 is a close analog of BI-3812 which binds only very weakly to the BCL6 BTB domain ($IC_{50} \sim 10 \mu M$).




Figure 4: BI-5273 which serves as a negative control

Selectivity

The intracellular selectivity profile was determined for the close analog BI-3802. For BI-3802 BCL6 was confirmed as the major target of this compound in DLBCL cells¹.

Co-crystal structure of the BI probe compound and the target protein.

Not available. However, the X-ray structure with the close analog BI-3802 was solved at Boehringer Ingelheim.

Reference molecule(s)

Several small molecule BCL6 inhibitors have been published recently^{7,8,9,10}.

Summary

BI-3812 is a single digit nanomolar BCL6::Co-repressor inhibitor which also inhibits the BCL6::Co-repressor complex formation in cells ($IC_{50} = 40$ nM). BI-3812 is a classical PPI inhibitor probe compound for testing hypotheses around BCL6 biology *in vitro*. With the close analogs BI-5273 as negative control and BI-3802 as BCL6 degrading inhibitor we also offer complementary tools.

References

1. N. Kerres *et al.* Chemically induced degradation of the oncogenic transcription factor BCL6 *Cell Rep.* **2017**, 20(12), 2860-2875. [DOI: 10.1016/j.celrep.2017.08.081](https://doi.org/10.1016/j.celrep.2017.08.081), [PubMed](#).
2. S. Zollman *et al.* The BTB domain, found primarily in zinc finger proteins, defines an evolutionarily conserved family that includes several developmentally regulated genes in *Drosophila* *Proc. Natl. Acad. Sci. U S A* **1994**, 91, 10717-10721. [PMC](#), [PubMed](#).
3. A.L. Dent *et al.* Control of inflammation, cytokine expression, and germinal center formation by BCL-6, *Science* **1997**, 276, 589-592. [DOI: 10.1126/science.276.5312.589](https://doi.org/10.1126/science.276.5312.589), [PubMed](#).
4. K. Basso *et al.*, Integrated biochemical and computational approach identifies BCL6 direct target genes controlling multiple pathways in normal germinal center B cells *Blood* **2010**, 115, 975-984. [DOI: 10.1182/blood-2009-06-227017](https://doi.org/10.1182/blood-2009-06-227017), [PubMed](#).
5. K. Basso and R. Dalla-Favera, Roles of BCL6 in normal and transformed germinal center B cells *Immunol. Rev.* **2012**, 247, 172-183. [DOI: 10.1111/j.1600-065X.2012.01112.x](https://doi.org/10.1111/j.1600-065X.2012.01112.x), [PubMed](#).
6. K. Hatzi and A. Melnick, Breaking bad in the germinal center: how deregulation of BCL6 contributes to lymphomagenesis *Trends Mol. Med.* **2014**, 20, 343-352. [DOI: 10.1016/j.molmed.2014.03.001](https://doi.org/10.1016/j.molmed.2014.03.001), [PubMed](#).
7. M. G. Cardenas *et al.* Rationally designed BCL6 inhibitors target activated B cell diffuse large B cell lymphoma *J. Clin. Invest.* **2016**, 126, 3351-3362. [DOI: 10.1172/JCI85795](https://doi.org/10.1172/JCI85795), [PubMed](#).
8. W. McCoull *et al.* Discovery of Pyrazolo[1,5-a]pyrimidine B-Cell Lymphoma 6 (BCL6) Binders and Optimization to High Affinity Macrocyclic Inhibitors, *J. Med. Chem.* **2017**, 60, 4386-4402. [DOI: 10.1021/acs.jmedchem.7b00359](https://doi.org/10.1021/acs.jmedchem.7b00359), [PubMed](#).
9. Y. Kamada *et al.* Discovery of a B-cell lymphoma 6 Protein-Protein Interaction Inhibitor by a Biophysics-driven Fragment-based Approach, *J. Med. Chem.* **2017**, 60, 4358-4368. [DOI: 10.1021/acs.jmedchem.7b00313](https://doi.org/10.1021/acs.jmedchem.7b00313), [PubMed](#).
10. T. Yasui *et al.* Discovery of a novel B-cell lymphoma 6 (BCL6)-corepressor interaction inhibitor by utilizing structure-based drug design, *Bioorg. Med. Chem.* **2017**, 25, 4876-4886. [DOI: 10.1016/j.bmc.2017.07.037](https://doi.org/10.1016/j.bmc.2017.07.037), [PubMed](#).