

PTK2 proteolysis-targeting chimera | BI-0319

Table of contents

Summary	2
Chemical Structure	2
Highlights	3
Target information	3
In vitro activity	4
<i>In vitro</i> DMPK and CMC parameters	6
Negative control	6
Selectivity	7
Reference molecule(s)	9
Supplementary data	9
References	10

Summary

BI-0319 is a potent and selective PROTAC (proteolysis-targeting chimera) aimed at triggering the intracellular destruction of the PTK2 protein.

Chemical Structure




Figure 1: 2-D structure of BI-0319, a highly selective and potent PTK2 PROTAC




Figure 2: 3-D structure of BI-0319, a highly selective and potent PTK2 PROTAC

Highlights

BI-0319 is a potent and selective PROTAC degrader of PTK2 kinase. This compound is a first-in-class low molecular weight degrader that tethers a PTK2 inhibitor to a CRL2VHL E3 ubiquitin ligase ligand to trigger the intracellular destruction of PTK2 in a reversible but long-lasting manner. This compound is suitable for *in vitro* studies.

Target information

Focal adhesion tyrosine kinase (PTK2) is a cytoplasmic protein tyrosine kinase that is overexpressed and activated in many types of advanced-stage solid cancers. PTK2 has been shown to play an important role in adhesion, spreading, motility, invasion, metastasis, survival, angiogenesis, epithelial to mesenchymal transition (EMT), cancer stem cells and the tumor microenvironment.^{2,3} Overexpression and activation of PTK2 is associated with several human malignant diseases, and is correlated with poor overall patient survival.⁴⁻⁶ The focal adhesion tyrosine kinase (PTK2) is often over-expressed in human hepatocellular carcinoma (HCC) and several reports have linked PTK2 depletion and/or pharmacological inhibition to reduced tumorigenicity.^{7,8} However, the clinical relevance of targeting PTK2 remains to be proven. Traditionally small molecules have been used to inhibit the action of a target protein by occupying and blocking a functional region of the protein. However, the disconnect between modulation of intracellular PTK2 autophosphorylation and growth inhibition as well as the often suboptimal selectivity profile of the inhibitors used makes it difficult to link the reported blockade of HCC tumour initiation and maintenance to PTK2 inhibition.

An alternative innovative approach is the development of proteolysis targeting chimeras (PROTACs), i.e. hetero bifunctional compounds consisting of one moiety that binds a Cullin RING E3 ubiquitin ligase linked to another that binds a desired protein of interest (POI), bringing the ligase and the POI into close spatial proximity. This hijacks the intrinsic catalytic activity of the E3 ligase and directs it toward the POI as a neo-substrate, triggering its poly-ubiquitination and subsequent proteasome-dependent degradation. As a result, a PROTAC acts as a degrader of the target as opposed to just an inhibitor, enabling the effective post-translational elimination of a target gene product in living organisms.⁹ This approach presents many advantages compared to conventional target inhibition. One of the most attractive features of the approach is that a PROTAC molecule acts sub-stoichiometrically, i.e. it only needs to bind a molecule of target once to induce its degradation, and then is released and set free to bind another molecule of target and carry on, as in a catalytic cycle. For this reason, the concentrations required for PROTACs to be active in cells tend to be much lower compared to those needed to be reached and maintained with inhibitors, which can lead to fewer off-target effects and a more selective chemical intervention on the desired target.




Figure 3: Structure of BI-4464 (ligand in BI-0319) bound to PTK2 (PDB ID 6I8Z). Hydrogen bonds to cysteine 502 and aspartic acid 564 are depicted in blue.

In vitro activity

The binary affinities for BI-0319 to PTK2 and to the VCB complex CRL2^{VHL} are 19 and 114 nM respectively. BI-0319 degrades the PTK2 protein in A549 cells with a DC₅₀ of 243 nM to about 80% (Table 1). BI-0319 (VHL-based) degrades PTK2 with a median DC₅₀ of 82 nM to > 80 % across a panel of eleven HCC cell lines (Table 2).

Table 1. Binary affinities of BI-4464 and BI-0319 for PTK2 and the respective PTK2 degradation data in A549 cells.

	BI-4463 (PKT2 INHIBITOR)	BI-0319 (PTK2 PROTAC)
E3 ligase ligand	-	POMA
PTK2 pIC ₅₀ *	7.8 ± 0.1	7.7 ± 0.1
CRBN-DDB1 TR-FRET pIC ₅₀	<4	6.9

A549 cells, 18h, pDC ₅₀ **	<4	6.7 ± 0.1
A549 cells, 18h, D _{max} [%]**	-	80 ± 9

*Thermo Fisher SelectScreen Kinase Profiling Services, Z'-Light, ATP@Km, pIC₅₀ ± STDEV

** Degradation activity is reported as concentration needed to achieve 50 % PTK2 protein degradation (pDC₅₀ ± STDEV) and maximal achievable protein degradation (D_{max}) relative to DMSO. PTK2 levels were determined by protein capillary electrophoresis and normalized to GAPDH. (N = 3)

Table 2. Degradation characteristics and effect on proliferation of BI-0319 and BI-4464 in HCC lines.

CELL LINE	BI-4463 (PKT2 INHIBITOR)		BI-0319 (PKT2 PROTAC)	
	pDC ₅₀	D _{max} [%]	pIC ₅₀ (proliferation)	pIC ₅₀ (proliferation)
SNU-387	7.1	79.0	4.7	5.4
HUH-1	6.2	59.0	5.2	5.3
Hep3B2.1-7	6.2	65.0	<4.6	5.5
HepG2	6.9	87.0	5.1	5.2
SK-Hep-1	7.4	66.0	<4.6	5.4
HLF	8.0	99.0	5.1	5.2
SNU-398	6.9	86.0	5.2	5.2
HUCCT1	7.6	85.0	5.0	5.1
HLE	7.0	79.0	5.4	5.4
HuH-7	7.1	87.0	5.2	5.4
SNU-423	7.1	79.0	4.7	5.4

*Please note that data are reported as pIC₅₀ which is calculated from IC₅₀ and therefore have no unit.

In vitro DMPK and CMC parameters

BI-0319 is poorly soluble at physiological pH, and is a PGP substrate (high Caco2 efflux ratio). Of note, the CRBN-based PROTAC BI-3663 was considerably less stable in cell assay buffer containing 10% FCS (M+18 and +32 observed) than the VHL-based PROTAC **BI-0319** (Table 3).

PROBE NAME / NEGATIVE CONTROL	BI-0319	BI-4206
MW [Da]	1061	1061
Solubility @ pH 6.8 [$\mu\text{g/ml}$]	<1	<1
CACO permeability @ pH 7.4 [$*10^{-6}$ cm/s]	<0.7	0.4
CACO efflux ratio	>15	61
Microsomal stability (human/mouse/rat) [% Q _H]	>95/>95/78	>95/90/54
Half-life (EMEM, 10% FCS) [h]	253	-
Plasma protein binding (human/mouse/rat/10% FCS) [%]	>99.2/>99.9/>99.8/96.0	>99.8/>99.9/>99.8/-

Negative control

cisVHL, BI-4206 is the (S) hydroxy diastereoisomer of BI-0319 and serves as a negative control for BI-0319 (Figure 3). While exhibiting BI-0319 comparable PTK2 binding affinity it no longer is able to bind and recruit VCB and therefore is not degrading PTK2 proteins in cells.




Figure 3: cisVHL, BI-4206 which serves as a negative control

Selectivity

In a 397 kinase panel BI-0319 beside PTK2 only inhibited LRRK2 and FES by more than 50% at 1 μ M (Figure 4). Interestingly, BI-0319 is more selective than the already highly selective PTK2 TKI BI-4464. We assume that the kinase selectivity panel for BI-3663 is comparable to BI-0319 since both exit vector and linker are identical for a distance of nine atoms from the piperidine moiety of the PTK2 ligand




Figure 4. Kinase selectivity panels of the PTK2 ligand BI-4464 and VHL PROTAC BI-0319. Bars indicate percent inhibition at 1 μ M of either compound.

Multiplexed isobaric tagging mass spectrometry was employed to assess the cellular selectivity of BI-0319 for PTK2 degradation and identify potential degradation off-targets in a quantitative and unbiased manner. Amongst the 6,008 proteins quantified in this analysis in A549 cells, PTK2 showed a distinct and significant change in abundance upon treatment with BI-0319. (Figure 5). BI-0319 did not induce any significant changes in abundance of other detectable kinases, thus confirming the high selectivity of both degraders within the kinase family. Of note, the two most prominent kinase off-targets of the inhibitor (LRRK2 and FES) were not detected in this dataset.

Interestingly BI-0319 also induced a significant change of PDE6D levels (Figure 5), a finding corroborated by immunoblot in A549 cells.





Figure 5. Total proteome analysis of A549 cells treated with BI-3019 for 18 h and compared to DMSO controls. Samples were run in biological triplicates and analyzed by mass spectrometry. Volcano plot displays \log_2 of fold-change in abundance versus $-\log_{10}$ of adjusted p value (N = 3).

SELECTIVITY DATA AVAILABLE	BI-0319	BI-4206
SafetyScreen44™ with kind support of 	Yes	Yes
Invitrogen®	Yes	No
DiscoverX®	No	No
Dundee	No	No

The PROTAC BI-0319 was tested by the Eric Fisher Laboratory - Dana-Farber Cancer Institute as part of their Degradation Proteomics Initiative.^{10,11} It induces selective degradation of PTK2 after 5 h of treatment at 3 μ M in the neuroblastoma cell line Kelly Cells.



Global protein quantification was used to explore the unbiased proteome-wide selectivity of BI-0319 induced degradation. Whole cell protein quantification was performed using label free quantification with the Fischer lab's diaPASEF workflow. Of the 7,742 proteins quantified in this experiment, only PTK2 was found to be significantly downregulated in response to BI-0319 treatment. Statistical analysis was performed using a moderated t-test in Bioconductor's limma package to generate hit lists containing log2 Fold Change and P-values for each protein. The data are also displayed in the scatterplot above.

Reference molecule(s)

There are currently no PTK2 PROTACs with the benchmarked selectivity.

Supplementary data

2-D structure files can be downloaded free of charge from [openMe](https://openme.org).

References

1. Popow J., Arnhof H., Bader G., Berger H., Ciulli A., Covini D., Dank C., Gmaschitz T., Greb P., Karolyi-Ozguer J., Koegl M., McConnell D. B., Pearson M., Rieger M., Rinnenthal J., Roessler V., Schrenk A., Spina M., Steurer S., Trainor N., Traxler E., Wieshofer C., Zoephel A., Ettmayer P. Highly Selective PTK2 Proteolysis Targeting Chimeras to Probe Focal Adhesion Kinase Scaffolding Functions. *J. Med. Chem.* **2019**, *62*, 2508-2520. [DOI: 10.1021/acs.jmedchem.8b01826](https://doi.org/10.1021/acs.jmedchem.8b01826), [PubMed](#).
2. Golubovskaya V. M. Targeting FAK in human cancer: from finding to first clinical trials. *Front Biosci (Landmark Ed)* **2014**, *19*, 687-706. [PMCID: PMC3952878](https://pubmed.ncbi.nlm.nih.gov/23952878/), [PubMed](#).
3. Sulzmaier F. J., Jean C., Schlaepfer D. D. FAK in cancer: mechanistic findings and clinical applications. *Nat Rev Cancer* **2014**, *14*, 598-610. [DOI: 10.1038/nrc3792](https://doi.org/10.1038/nrc3792), [PubMed](#).
4. Judson P. L., He X., Cance W. G., Van Le L. Overexpression of focal adhesion kinase, a protein tyrosine kinase, in ovarian carcinoma. *Cancer* **1999**, *86*, 1551-1556. [DOI: 10.1002/\(sici\)1097-0142\(19991015\)86:6<1551::aid-cnrc23>3.0.co;2-p](https://doi.org/10.1002/(sici)1097-0142(19991015)86:6<1551::aid-cnrc23>3.0.co;2-p), [PubMed](#).
5. Lark A. L., Livasy C. A., Calvo B., Caskey L., Moore D. T., Yang X., Cance W. G. Overexpression of focal adhesion kinase in primary colorectal carcinomas and colorectal liver metastases: immunohistochemistry and real-time PCR analyses. *Clin Cancer Res* **2003**, *9*, 215-222. [PMID: 12538472](https://pubmed.ncbi.nlm.nih.gov/12538472/), [PubMed](#).
6. Miyazaki T., Kato H., Nakajima M., Sohda M., Fukai Y., Masuda N., Manda R., Fukuchi M., Tsukada K., Kuwano H. FAK overexpression is correlated with tumour invasiveness and lymph node metastasis in oesophageal squamous cell carcinoma. *Br J Cancer* **2003**, *89*, 140-145. [doi: 10.1038/sj.bjc.6601050](https://doi.org/10.1038/sj.bjc.6601050), [PubMed](#).
7. Li W. X., Chen L. P., Sun M. Y., Li J. T., Liu H. Z., Zhu W. 3'-Diindolylmethane inhibits migration, invasion and metastasis of hepatocellular carcinoma by suppressing FAK signaling. *Oncotarget* **2015**, *6*, 23776-23792. [DOI: 10.18632/oncotarget.4196](https://doi.org/10.18632/oncotarget.4196), [PubMed](#).
8. Gnani D., Romito I., Artuso S., Chierici M., De Stefanis C., Panera N., Crudele A., Ceccarelli S., Carcarino E., D'Oria V., Porru M., Giorda E., Ferrari K., Miele L., Villa E., Balsano C., Pasini D., Furlanello C., Locatelli F., Nobili V., Rota R., Leonetti C., Alisi A. Focal adhesion kinase depletion reduces human hepatocellular carcinoma growth by repressing enhancer of zeste homolog 2. *Cell Death Differ* **2017**, *24*, 889-902. [DOI: 10.1038/cdd.2017.34](https://doi.org/10.1038/cdd.2017.34), [PubMed](#).
9. Bondeson D. P., Mares A., Smith I. E., Ko E., Campos S., Miah A. H., Mulholland K. E., Routly N., Buckley D. L., Gustafson J. L., Zinn N., Grandi P., Shimamura S., Bergamini G., Faeltsh-Savitski M., Bantscheff M., Cox C., Gordon D. A., Willard R. R., Flanagan J. J., Casillas L. N., Votta B. J., den Besten W., Famm K., Kruidenier L., Carter P. S., Harling J. D., Churcher I., Crews C. M. Catalytic in vivo protein knockdown by small-molecule PROTACs. *Nat Chem Biol* **2015**, *11*, 611-617. [DOI: 10.1038/nchembio.1858](https://doi.org/10.1038/nchembio.1858), [PubMed](#).
10. The data can be accessed via the following link: [Proteomics Dashboard \(fischerlab.org\)](https://proteomics.fischerlab.org/)

11. Donovan K. A., Ferguson F. M., Bushman J. W., Eleuteri N. A., Bhunia D., Ryu S., Tan L., Shi K., Yue H., Liu X., Dobrovolsky D., Jiang B., Wang J., Hao M., You I., Teng M., Liang Y., Hatcher J., Li Z., Manz T. D., Groendyke B., Hu W., Nam Y., Sengupta S., Cho H., Shin I., Agius M. P., Ghobrial I. M., Ma M. W., Che J., Buhrlage S. J., Sim T., Gray N. S., Fischer E. S. Mapping the Degradable Kinome Provides a Resource for Expedited Degradation Development *Cell*. **2020**, 183(6):1714-1731. [DOI: 10.1016/j.cell.2020.10.038](https://doi.org/10.1016/j.cell.2020.10.038), [PubMed](#).